

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 0 645 145 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention  
of the grant of the patent:  
12.03.1997 Bulletin 1997/11

(51) Int. Cl.<sup>6</sup>: **A61K 47/44**, A61K 47/10

(21) Application number: **94115180.5**

(22) Date of filing: **27.09.1994**

(54) **Stabilized pharmaceutical composition and stabilizing solvent**

Stabilisierte Arzneimittel und stabilisierendes Lösungsmittel  
Composition pharmaceutique stabilisée et solvant stabilisant

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL  
PT SE**

(30) Priority: **29.09.1993 US 128026**

(43) Date of publication of application:  
**29.03.1995 Bulletin 1995/13**

(73) Proprietor: **BRISTOL-MYERS SQUIBB COMPANY**  
**Princeton, NJ 08543-4000 (US)**

(72) Inventors:  
• **Agharkar, Shreerham Narahari**  
**Fayetteville, NY 13066 (US)**  
• **Gogate, Uday S.**  
**East Syracuse, NY 13057 (US)**  
• **Cavanak, Thomas**  
**CH-4105 Biel-Benken (CH)**

(74) Representative: **Josif, Albert, Dr.-Ing. et al**  
**Baaderstrasse 3**  
**80469 München (DE)**

(56) References cited:  
**WO-A-94/12030** **GB-A- 2 221 157**

- **CHEMICAL ABSTRACTS**, vol. 109, no. 14, 3 October 1988, Columbus, Ohio, US; abstract no. 111136, 'PURIFICATION OF POLYETHYLENE GLYCOL FOR USE IN PHARMACEUTICALS' & RO-A-92 831 (COMBINATUL PETROCHIMIC) 31 December 1987
- **PATENT ABSTRACTS OF JAPAN** vol. 010, no. 350 (C-387) 26 November 1986 & JP-A-61 152 621 (LION CORP.) 11 July 1986 & DATABASE WPI Week 8638, Derwent Publications Ltd., London, GB; AN 86-247656 & JP-A-61 152 621 (LION CORP.) 11 July 1986
- **DRUG DEVELOPMENT AND INDUSTRIAL PHARMACY**, vol.18, no.1, 1992 pages 1 - 19 M.S. PATEL ET AL 'FACTORS AFFECTING THE CHEMICAL STABILITY OF CARBOXYLIC ACID DRUGS IN ESS SOFTGEL FORMULATIONS BASED ON PEG'
- **PATENT ABSTRACTS OF JAPAN**, vol. 010, no. 350 (C-387), 26 November 1986; & JP-A-61 152 621

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 0 645 145 B1**

**Description**Field of the Invention

5 The present invention is directed to a solvent system suitable for preparing a stabilized injection concentrate containing a pharmaceutical agent. More particularly, the invention is directed to a stabilized injection concentrate using a treated solvent system having a reduced carboxylate anion content and a method for stabilizing a pharmaceutical composition.

Background of the Invention

Pharmaceutical compositions usually require a suitable solvent or carrier system to disperse the active agent to enable the composition to be administered to a patient. The solvent must typically be capable of solubilizing or dispersing a therapeutically effective amount of the active agent to produce an effective injection composition. Moreover, the solvent system must be compatible with the active agent and be non-toxic to the patient.

15 Numerous pharmaceutical agents are not sufficiently soluble in any one solvent to enable the resulting composition to be effective. To overcome the disadvantages of the limitations of the solvent to solubilize the active agent, mixtures of two or more solvents are sometimes used. These co-solvent systems are suitable for solubilizing many pharmaceutical agents which cannot otherwise be solubilized or dispersed in a carrier system.

20 One example of a co-solvent system incorporates a mixture of a polar solvent and a non-ionic solvent, such as a mixture of a polyethylene glycol and Cremophor EL. Cremophor EL is a condensation product of castor oil and ethylene oxide sold by BASF. Another suitable co-solvent system for many pharmaceutical agents is a 50:50 mixture of ethanol and Cremophor EL. Although these co-solvent systems can be effective in solubilizing many compounds, they are not without their disadvantages. For example, co-solvents of ethanol and Cremophor are known to result in particulates forming upon dilution with infusion solutions. In addition, fibrous precipitates of unknown composition form in some compositions during storage for extended periods of time. It is generally believed that the precipitates are decomposition byproducts of either components in the solvent or the solubilized agent.

30 In WO91/02531, published March 7, 1991, Cremophor is disclosed as reversing the multi-drug resistance phenotype of a tumor cell without altering the drug sensitivity of the parent cell line. Cremophor is also disclosed to increase hemopoiesis reconstituting capacity and/or maintain hemopoiesis reconstituting capacity following perturbation of bone marrow to protect a patient during radiation and/or chemotherapy cancer treatments.

Another example of a pharmaceutical composition including a co-solvent system is Taxol which contains paclitaxel in a 50:50 mixture of ethanol and Cremophor EL. Paclitaxel isolated from the Pacific yew bark which has been used to treat cancer patients. Although the ethanol and Cremophor EL co-solvent system is effective in solubilizing sufficient amounts of the paclitaxel, the resulting composition has been shown to have a limited shelf life. During storage for extended periods of time, the potency or pharmaceutical activity of the composition can decrease as much as 60%.

35 It has been discovered that commercial grade Cremophor EL with ethanol as a co-solvent, although effective in solubilizing pharmaceutical agents, produces injection compositions that exhibit instability over extended periods of time. In particular, pharmaceutical compositions of Taxol in a co-solvent of 50:50 by volume of dehydrated ethyl alcohol and commercial grade Cremophor EL exhibit a loss of potency of greater than 60% after storage for 12 weeks at 50°C. The loss of potency is attributed to the decomposition of paclitaxel during storage.

40 Previous efforts to develop a shelf stable composition of some pharmaceutical compositions in various co-solvent systems have not been entirely successful. Thus, there is a continuing need in the art for a co-solvent system capable of being used for preparing stabilized compositions and, in particular, stabilized injection compositions containing a pharmaceutical agent.

45 WO-A-94/12030 which is considered prior art under the terms of Article 54(3) and (4) EPC discloses compositions comprising paclitaxel, polyethoxylated castor oil and ethanol adjusted to a pH less than 8.1 by addition of an acid.

Summary of the Invention

50 The invention is directed to a solvent system and in particular a co-solvent system suitable for preparing stabilized injection compositions containing at least one pharmaceutical agent. Accordingly, it is a primary object of the invention to provide a method of preparing stabilized pharmaceutical compositions including the novel solvent system.

55 The stabilized pharmaceutical composition produced from the solvent system of the invention has been shown to have a shelf life greater than the previous compositions. The co-solvent system of the invention is particularly suitable for use with pharmaceutical compounds that exhibit decomposition which is catalyzed by carboxylate anions. Of interest are the antineoplastic agents paclitaxel, camptothecin and derivatives thereof.

The stabilized composition of the invention comprises at least one pharmaceutical compound selected from the group consisting of paclitaxel, camptothecin and derivatives thereof; and a solvent capable of dispersing or solubilizing

said pharmaceutical compound, said solvent comprising an effective amount of a non-ionic solubilizing agent having a carboxylate anion content sufficiently low to prevent catalyzed degradation of said pharmaceutical compound, provided that compositions comprising paclitaxel, polyethoxylated castor oil and ethanol adjusted to a pH less than 8.1 by addition of an acid other than HBr, HF and HI are excluded.

The solvent system of the invention includes a non-ionic solubilizing agent. The solvent system typically includes a solvent and a solubilizing agent. In preferred forms of the invention, the solubilizing agents are polyoxyalkylene modified lipids and alkylene oxides such as polyethylene glycol and derivatives thereof. The solubilizing agent can be a condensation product of an alkylene oxide and a lipid or fatty acid. The preferred solvent system includes a polyoxyethylated castor oil such as that sold under the tradename Cremophor EL. The Cremophor EL is treated to reduce the carboxylate anion content to a sufficiently low concentration to minimize the decomposition of the pharmaceutical agent that is catalyzed by the carboxylate anions. The carboxylate anion content of the Cremophor EL is lowered by either contacting the Cremophor EL with an aluminum oxide bed to separate the carboxylate anions as well as other impurities or by the addition of an acid and particularly a mineral acid such as HCl or HNO<sub>3</sub>. In further embodiments, the solvent is treated with a reactant to reduce the carboxylate anion or convert the carboxylate anion to a non-reactive form.

Further advantages of the invention are attained by providing a method of stabilizing a pharmaceutical composition containing a pharmaceutical agent selected from the group consisting of paclitaxel, camptothecin and derivatives thereof and a solvent containing ethanol and a solubilizing amount of at least one solubilizing agent as defined in claim 1, the method comprising treating the solvent to reduce the carboxylate content to a sufficiently low level to substantially prevent carboxylate anion catalyzed degradation of the pharmaceutical agent.

#### Detailed Description of the Invention

The disadvantages and limitations of the previous injection composition and solvent systems are overcome by the present invention while providing a convenient and efficient method of stabilizing pharmaceutical compositions suitable for injection.

It has been surprisingly discovered that paclitaxel reacts with ethanol during storage and that the decomposition of paclitaxel is catalyzed by the carboxylate anions in the solvent. Lowering the carboxylate concentration of the solvent has been found to produce a stabilizing effect in the pharmaceutical composition. The lower carboxylate concentration extends the shelf life of the composition by reducing the rate of decomposition of the pharmaceutical agent and reducing the formation of decomposition byproducts.

The solvent system has a sufficiently low impurity content to provide a stabilizing effect in pharmaceutical compositions. The resulting pharmaceutical composition produced by the method can be used to prepare injection compositions by diluting with a pharmaceutically acceptable carrier or diluent. The composition is sufficiently stable to minimize degradation of the active compound and to reduce the loss of potency during storage. The diluted composition does not exhibit the formation of precipitates so that the pharmaceutically active agent can be stored in a readily usable form.




The pharmaceutical composition includes at least one pharmaceutical compound having a tendency to degrade during storage where the decomposition reaction is catalyzed by carboxylate anions. The pharmaceutical compounds have antineoplastic activity such as, for example, paclitaxel sold under the tradename Taxol. The various analogs and derivatives of Taxol may also be used.

Taxol is prepared as an injection concentrate containing 6 mg/ml paclitaxel in 50:50 by volume ethanol and polyoxyethylated castor oil. The polyoxyethylated castor oil is sold under the tradename Cremophor EL. Paclitaxel is an active component isolated from the bark of Pacific yew trees. Recently, paclitaxel has been found to be produced in small amounts from a fungus found in the bark of yew trees. Paclitaxel is known to have antineoplastic activity.

Another suitable pharmaceutical agent is Camptothecin having the chemical name 4-ethyl-hydroxy-1H-pyrano-[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H, 12H)dione. Camptothecin is isolated from the stem wood of the Chinese tree and has been shown to exhibit antileukemic and antitumor activity.

The pharmaceutical agents paclitaxel and camptothecin exhibit degradation and loss of activity during storage. The solvent system and process of the invention are particularly advantageous for use with these pharmaceutical agents that react with or are unstable in the solvent. In particular, the pharmaceutical agents of interest have an ester linkage that can be cleaved by an alcohol in the presence of carboxylate anions.

Paclitaxel is typically produced as a concentrate or solution in a vehicle suitable for injection in the amount of 6 mg/ml. The vehicle is usually a mixture of ethanol and Cremophor EL in the amount of 50:50 by volume. During storage, the activity of the paclitaxel is known to decrease. Paclitaxel has the Formula I, which HPLC shows to degrade into Baccatin III of Formula II and the ethyl ester side chain of Formula III.



55 The solvent is a co-solvent mixture of at least one solvent and a non-ionic solubilizing agent. The preferred solvents include the alcohols such as dehydrated ethanol and the pharmaceutically acceptable polyols such as polyethyleneglycol. The solubilizing agent in preferred embodiments is a polyoxyethylated castor oil such as that sold under the trade-name Cremophor EL by BASF. The co-solvent in preferred embodiments contains about 40% to 60% by volume of the polyoxyethylated castor oil with the balance being an alcohol or polyol. In a particularly preferred embodiment of the

invention, the co-solvent comprises about 50:50 by volume of dehydrated ethanol and Cremophor EL.

The co-solvent of the invention preferably includes a non-ionic surfactant as the solubilizing agent with Cremophor EL being the most preferred. Cremophor EL is a condensation product of castor oil and about 20 to 40 moles and preferably 30 to 35 moles of ethylene oxide per mole of castor oil. Cremophor EL may be prepared by the method disclosed in U.S. Patent No. 3,070,499. Cremophor EL is also known by its common names polyoxyethylene-glycerol triricinoleate and glycerol-polyethyleneglycol ricinoleate. The biological and chemical equivalents or derivatives of Cremophor EL may also be used.

In alternative embodiments, the non-ionic surfactant or solubilizing agent may include other ethylene oxide modified lipids, hydroxylated tallow oils, polysorbate 80, also known as Tween 80, polyethoxylated 12-hydroxy stearic acid, polyoxyethylene sorbitan fatty acid esters, polyethylene glycol esters, polyethylene fatty acid esters, block co-polymers of ethylene oxide and propylene oxide, ethylated fatty alcohol ethers, and octyl-phenoxy polyethoxy ethanol compounds. These non-ionic, solubilizing agents can be produced by methods well known in the art or from commercial suppliers. The solubilizer may include other condensation products of alkylene oxides, although the alkylene oxide modified lipids are generally preferred. Examples of suitable solubilizing agents are PEG 400 and PEG 40 castor oil.

The carboxylate anion content of the solvent can be lowered by a number of methods. In a first embodiment of the invention, the Cremophor EL or other solvent is passed through a standard chromatography column of aluminum oxide. The aluminum oxide adsorbs the carboxylate anions as well as other impurities to reduce the carboxylate anion content of the solvent.

In an alternative embodiment of the invention, the solvent is treated by the addition of an acid in a stabilizing amount to reduce the carboxylate anion content to a sufficiently low level to substantially prevent catalyzed degradation of the pharmaceutical compound, provided that the method excludes the addition of acid other than HBr, HF and HI to lower the pH of a composition comprising paclitaxel, polyethoxylated castor oil and ethanol to below 8.1. The acid may be added to the solvent before or after admixing with the pharmaceutical compound. Generally, mineral acids such as, for example, HCl, HBr, HF, HI,  $\text{H}_2\text{SO}_4$  and  $\text{HNO}_3$  are used. Alternatively, organic acids such as acetic may be used. The organic acids are generally less preferred since they can provide a source of carboxylate anions to hinder the stabilizing effect of the acid treatment. The acid is preferably added in amounts to provide  $5.6 \times 10^{-6}$  to  $8.4 \times 10^{-6}$  grams of  $\text{H}^+$  per ml of the solvent. Effective stabilization of the composition is obtained by the addition of the acid to provide about  $7.0 \times 10^{-6}$  grams of  $\text{H}^+$  per ml. The acid is added to lower the carboxylate anion content to less than or equal to about  $0.6 \times 10^{-6}$  gram equivalents of carboxylate anion per ml of solvent.

Taxol solutions containing  $0.6 \times 10^{-6}$  gram equivalents of carboxylate anion have exhibited 94% paclitaxel remaining after storage at  $50^\circ\text{C}$  for 28 days. 1.0 gram equivalents of anions is neutralized by 1.0 gram  $\text{H}^+$  per equivalent.

The amount of acid added to the solvent will depend on the amount of the non-ionic surfactant in the co-solvent system and the type of the surfactant. The non-ionic surfactants which by their method of preparation inherently contain greater amounts of carboxylate anion impurities will typically require larger amounts of acid to lower the carboxylate anion content.

In further embodiments the solvent may be treated with other reactants which are capable of lowering the carboxylate anion concentration of the solution. For example, insoluble salts of the carboxylates or derivatives of carboxylates may be formed. It is generally preferred to remove the carboxylate anions using a reagent which does not form precipitates in the composition. The addition of a strong mineral acid is generally preferred since the mineral acids and the resulting carboxylic acids are non-toxic and readily solubilized in the solvent. When the carboxylate anions are reacted in a manner which form an insoluble precipitate, the precipitate should be separated from the composition prior to administering the composition to a patient.

The following examples are intended to demonstrate the preferred embodiments of the invention. One skilled in the art will readily recognize that numerous embodiments of the invention can be practiced to achieve the stabilizing effect.

#### EXAMPLE 1

This example analyzes the properties of Taxol and paclitaxel in a solution of ethanol and Cremophor EL. The Taxol was obtained containing 6 mg/ml of paclitaxel in a 50:50 mixture by volume of Cremophor EL and dehydrated ethanol. A first group of samples 1-4 were prepared using different batches of processed Cremophor EL and a second group of samples 5-7 were prepared using different batches of un-processed commercial grade Cremophor EL. The processed Cremophor EL was prepared by passing 100 Kg of the Cremophor EL through a standard chromatography column containing 19.5 Kg of aluminum oxide sold under the tradename CAMAG. In order to understand the cause of the Taxol instability in unprocessed Cremophor EL, each sample was analyzed by observing the Karl-Fischer moisture, potassium ion content, acid value, and peroxide value. The results of these observations are presented in Table 1.

TABLE 1

Type of Cremophor EL	Sample No.	KF Moisture %	Potassium level (ppm)	Acid Value	Peroxide Value
Processed	Sample 1	0.2	<1	0.2	8
	Sample 2	0.3	2	0.3	8
	Sample 3	0.4	2	0.2	9
	Sample 4	0.8	4	0.3	3
Un-Processed	Sample 5	2.7	463	3.4	18
	Sample 6	2.6	482	1.6	n.d.
	Sample 7	2.6	n.d.	1.4	15

n.d. = not determined

The data of Table 1 show consistency of moisture content, potassium content, acid value and peroxide value between the batches of processed Cremophor EL used to make samples 1-4. The data further shows consistently higher levels of moisture, potassium content, acid values and peroxide values for the unprocessed Cremophor EL.

#### EXAMPLE 2

This example was carried out to demonstrate the instability of Taxol and paclitaxel in a co-solvent containing Cremophor EL and to determine the nature of the decomposition products. Specifically, this example was carried out to determine whether high levels of moisture, potassium, acid value and peroxide value of unprocessed Cremophor EL as determined in Example 1 are related to the rate of decomposition and loss of potency of Taxol.

Samples 8-21 were prepared by dissolving 6 mg/ml of paclitaxel in 50:50 v/v mixture of processed Cremophor EL and dehydrated ethanol. The Cremophor EL of samples 8-21 was processed as discussed above in Example 1. Sample 22 was prepared as a control sample from unprocessed Cremophor EL in a 50:50 v/v mixture with paclitaxel in the amount of 6 mg/ml. Each of the samples 8-20 were mixed with the components listed in Tables 2 and 3. Three ml aliquots of the test samples were placed in 6 cc Type I flint glass vials. The vials were stoppered with West 4455/45 Teflon-faced stoppers, sealed and stored for four weeks at 50°C, and then analyzed by HPLC for Taxol concentration. The control parameters of the HPLC are as follows:

Column	Jones Cyano RP 5 $\mu$ , 25cm x 4.6mm ID
Detector Wavelength	227 nm
Mobile phase	35% acetonitrile:65% 20mM acetate buffer (pH 4)
Flow rate	1.5 mL/min.
Diluent	Mobile phase
Sample Concentration	0.05 mg/mL
Injection volume	20 microliters
Retention time	Taxol 10.5 mins.

The pH value for each of the samples was noted following 1:10 dilution with water. Peroxide value (EP method), acid value (USP method), moisture content and potassium levels were determined as shown in Tables 2 and 3.

TABLE 2

Sample No.	Component Added	% KF Moisture	Initial pH <sup>a</sup>	Following storage for 28 days at 50°C	
				% taxol remaining	pH <sup>a</sup>
Sample 8 <sup>b</sup>	4% H <sub>2</sub> O <sub>2</sub> solution	5.5	4.6	94.9	3.6
Sample 9 <sup>c</sup>	10% H <sub>2</sub> O <sub>2</sub> solution	9.6	4.6	96.8	3.5
Sample 10 <sup>d</sup>	4% water	5.8	5.0	94.5	4.2
Sample 11 <sup>e</sup>	Acetic acid (1.5 mg/mL)	2.6	3.8	94.1	4.1
Sample 12 <sup>f</sup>	None (control)	2.6	4.4	95.5	4.5

<sup>a</sup>Following 1:10 dilution with Water for Injection.<sup>b</sup>Peroxide value = 232.<sup>c</sup>Peroxide value = 652.<sup>d</sup>Peroxide value = 14.<sup>e</sup>Acid value = 1.2.<sup>f</sup>Acid value = 0.5, Peroxide value = 4.

TABLE 3

Sample No.	Component Added	K <sup>+</sup> level (ppm)	Initial pH <sup>a</sup>	Following Storage for 28 Days at 50°C.	
				% Taxol Remaining	pH <sup>a</sup>
Sample 13	CH <sub>3</sub> COOK 0.126% w/v	530	5.6	19.3	5.1
Sample 14	KCl 0.10% w/v	551	6.0	98.8	4.7
Sample 15	NaCl 0.125 %w/v	4	6.3	96.8	4.9
Sample 16	CH <sub>3</sub> COONH <sub>4</sub> 0.10% w/v	4	5.8	63.7	4.7
Sample 17	CH <sub>3</sub> COON <sub>a</sub> 0.104 %w/v	5	6.1	5.8	5.5
Sample 18	CH <sub>3</sub> CH <sub>2</sub> COOK 0.144 %w/v	468	5.4	15.2	5.1
Sample 19	Linolelaidic acid (LA) 0.2% W/V	5	6.4	99.5	5.4
Sample 20	K salt of LA 0.228% w/v	150	6.3	52.8	6.2
Sample 21	None (Control)	<2	4.4	95.5	4.5
Sample 22	prepared with unprocessed Cremophor EL	240	4.6	86.5	4.8

<sup>a</sup>Following 1:10 dilution with Water for Injection.

As shown in Table 2, the stability of Taxol as a solution of a 50:50 mixture of dehydrated ethanol and Cremophor EL is not affected by the addition of hydrogen peroxide, water, or acetic acid, thereby demonstrating that the Taxol instability is not related to peroxide value, moisture content or acid value of the Cremophor EL. Each of samples 8-11 exhibited comparable loss of Taxol potency regardless of the reagent added. In each of samples 8-11, the percent of Taxol remaining reflecting the amount of paclitaxel in the samples ranged from 94.1% to 96.8% compared to 95.5% of the control Sample 12.

Sample 13 of Table 3 containing potassium acetate is shown to be less stable than control Sample 21 which contained no added component. Sample 14 containing potassium chloride is shown to be comparable with the control sample 21. These data suggest that the stability of the Taxol is not affected by the presence of potassium ions but rather are

## EP 0 645 145 B1

adversely affected by the anionic form of the carboxylic acids. Sample 13 containing potassium acetate, sample 16 containing ammonium acetate, sample 17 containing sodium acetate, sample 18 containing potassium propionate and Sample 20 containing potassium linoleate are all less stable than the control sample 22. However, the stability of sample 11 containing acetic acid and Sample 19 containing linoleic acid is comparable to control sample 22 indicating that the carboxylic acids in the non-ionized form do not affect Taxol stability. Sample 22 containing unprocessed Cremophor EL is also shown to be less stable than control Sample 21 containing processed Cremophor EL. HPLC of each of the samples indicated the degradation products to be Baccatin III and the ethyl ester of the Taxol side chain.

### EXAMPLE 3

This example is not an embodiment of the invention but contains information applicable to the invention.

This example demonstrates the stabilizing effect of acids added to Taxol samples containing unprocessed Cremophor EL. The samples in this example were prepared from Taxol containing 6 mg/ml paclitaxel in a 50:50 v/v mixture of dehydrated ethanol and unprocessed commercial grade Cremophor EL. As indicated in Table 4, samples 23-28 were mixed with HCl, sample 29 was mixed with acetic acid and Sample 30 was mixed with nitric acid. Sample 31 was the control sample containing no added acid. Sample 32 containing acetic acid and control sample 33 containing no added acid were also prepared from unprocessed Cremophor EL. The samples were separated into 3 ml aliquots and placed into 5 cc Type 1 flint glass vials and stoppered with Daikyo #713 fluoresein coated closures. The vials were stored at 50°C for 56 days and analyzed by HPLC for paclitaxel (Taxol) concentration. The HPLC parameters were as follows:

Column	Jones Cyano RP 5 $\mu$ , 25cm x 4.6mm ID
Detector Wavelength	227 nm
Mobile phase	35% acetonitrile:65% 20mM acetate buffer (pH 4)
Flow rate	1.5 mL/min.
Diluent	Mobile phase
Sample Concentration	~0.05 mg/mL
Injection volume	20 microliters
Retention time	paclitaxel ~ 10.5 mins.

The data in Table 4 demonstrate that the stabilizing effect of the acid increased with increasing amounts of the acid. These results are consistent with the premise that the degradation of Taxol is due to the presence of carboxylate anions rather than the acid. The data further demonstrates that the mineral acids such as HCl and HNO<sub>3</sub> are better stabilizing agents with greater than 97% of the Taxol remaining after 56 days when the acid is added at levels to provide 5.6 x 10<sup>-6</sup> to 8.4 x 10<sup>-6</sup> grams of H<sup>+</sup> per ml. The mineral acids showed a greater stabilizing effect than acetic acid even when the acetic acid was added at a higher level of 70 x 10<sup>-6</sup> grams of H<sup>+</sup> per ml. These results are due to the acid strength of the mineral acids compared to organic acids to neutralize the carboxylate anions. In addition, the acetic acid as a source of carboxylate anions can contribute to some paclitaxel degradation. The amount of paclitaxel remaining in samples 26 and 30 being greater than 100% is due to analytical variations in the measurements.



TABLE 4

TAXOL Injection Sample No.	Acid added (Gms of H <sup>+</sup> added per mL)	Initial pH <sup>c</sup>	Following Storage for 56 Days at 50° C.	
			% Paclitaxel Remaining	pH <sup>c</sup>
Sample 23 <sup>a</sup>	HCl (3.5x10 <sup>-6</sup> )	5.1	62.8	4.5
Sample 24 <sup>a</sup>	HCl (5.6x10 <sup>-6</sup> )	3.9	98.1	3.8
Sample 25 <sup>a</sup>	HCl (6.3x10 <sup>-6</sup> )	3.8	97.9	3.8
Sample 26 <sup>a</sup>	HCl (7.0x10 <sup>-6</sup> )	3.8	100.9	3.6
Sample 27 <sup>a</sup>	HCl (7.7x10 <sup>-6</sup> )	3.6	99.3	3.6
Sample 28 <sup>a</sup>	HCl (8.4x10 <sup>-6</sup> )	3.6	99.1	3.6
Sample 29 <sup>a</sup>	CH <sub>3</sub> COOH (7.0x10 <sup>-6</sup> )	4.4	69.5	4.5
Sample 30 <sup>a</sup>	HNO <sub>3</sub> (7.0x10 <sup>-6</sup> )	3.7	100.4	3.8
Sample 31 <sup>a</sup>	Control (No acid added)	5.5	49.0	5.0
Sample 32 <sup>b</sup>	CH <sub>3</sub> COOH (7.0x10 <sup>-6</sup> )	3.7	87.8	3.7
Sample 33 <sup>b</sup>	Control (No acid added)	6.3	22.6	5.5

<sup>a</sup>Unprocessed Cremophor EL, BASF Lot No. 98-2384, was used in these solutions.

<sup>b</sup>Unprocessed Cremophor EL, BASF Lot No. 14-1213, was used in these solutions.

<sup>c</sup>Following 1:10 dilution with Water for Injection.

Samples 34A, 34B, 35A, 35B, 36A and 36B as shown in Table 5 were prepared using different lots of Cremophor to show the consistency of the stabilizing effect of mineral acids. Each sample was prepared containing 6 mg/ml paclitaxel in a 50:50 mixture of dehydrated ethanol and unprocessed Cremophor EL. Samples 34A, 35A and 36A were prepared in the manner of Samples 23-28 and to contain HCl in an amount to provide  $7.0 \times 10^{-6}$  of H<sup>+</sup> per ml. Control Samples 34B, 35B and 36B did not contain added acid. The samples were stored in closed vials for 56 days at 50°C. The amount of paclitaxel remaining was determined by HPLC as shown in Table 5.

TABLE 5

Cremophor EL Lot No.-unprocessed	Acid added	Initial pH <sup>a</sup>	Following Storage for 56 Days at 50° C.	
			% Paclitaxel Remaining	pH <sup>a</sup>
Sample 34A	HCl	4.0	98.4	3.9
Sample 34B	Control <sup>b</sup>	6.1	25.0	5.4
Sample 35A	HCl	3.9	97.8	4.0
Sample 35B	Control <sup>b</sup>	6.3	22.6	5.5
Sample 36A	HCl	3.8	100.9	3.6
Sample 36B	Control <sup>b</sup>	5.5	49.0	5.0

<sup>a</sup>Following 1:10 dilution with Water for Injection.

<sup>b</sup>No acid was added in the control solutions.

The data of Table 5 demonstrate the consistent stabilizing effect of mineral acids with greater than 97% of the paclitaxel remaining after 56 days compared to less than 50% remaining in the control samples.

The data of Examples 2 and 3 demonstrate the stabilizing effect of processing the Cremophor EL by contacting with aluminum oxide or treating with an acid to reduce the carboxylate content of the solvent. It will be appreciated by those skilled in the art that the process of reducing the carboxylate anion level can be used with other solvents and solubilizers. Taxol has been demonstrated to exhibit increased shelf life from the stabilizing effect of the acids and the reduction in carboxylate anion content.

## Claims

1. A stabilized composition comprising at least one pharmaceutical compound selected from the group consisting of paclitaxel, camptothecin and derivatives thereof; and  
a solvent capable of dispersing or solubilizing said pharmaceutical compound, said solvent comprising an effective amount of a non-ionic solubilizing agent having a carboxylate anion content sufficiently low to prevent catalyzed degradation of said pharmaceutical compound,  
provided that compositions comprising paclitaxel, polyethoxylated castor oil and ethanol adjusted to a pH less than 8.1 by addition of an acid other than HBr, HF and HI are excluded.
2. A stabilized composition comprising at least one pharmaceutical compound selected from the group consisting of paclitaxel, camptothecin and derivatives thereof; and  
a solvent capable of dispersing or solubilizing said pharmaceutical compound, said solvent comprising an effective amount of a non-ionic solubilizing agent having a carboxylate anion content sufficiently low to prevent catalyzed degradation of said pharmaceutical compound wherein, prior to dispersing said pharmaceutical compound in said solvent, said solvent has been treated to reduce the carboxylate anion content, said treatment comprising contacting said solvent with aluminum oxide.
3. The composition of claim 1 or 2, wherein said solvent has a carboxylate anion content of about  $0.6 \times 10^{-6}$  gram equivalents of carboxylate anion per ml of solvent.
4. The composition of claim 1, 2 or 3, wherein said solvent comprises a condensation product of alkylene oxide and a lipid, more particularly polyoxyethylated castor oil or polyoxyethylated 12-hydroxy stearic acid.
5. The composition of claim 4, wherein said polyoxyethylated lipid condensation product is made of castor oil and 20 to 40 moles of ethylene oxide per mole of castor oil.
6. The composition of any one of claims 1 to 5, wherein said solvent further comprises an alcohol, more specifically an alcohol selected from the group consisting of ethanol and polyethylene glycol.
7. The composition of claim 6, wherein said solvent is a mixture of ethyl alcohol and said polyoxyethylated castor oil in a ratio of about 50:50 by volume.
8. The composition of any one of claims 1 to 7, wherein said solvent contains an effective amount of an acid to maintain a carboxylate anion content of less than or equal to about  $0.6 \times 10^{-6}$  gram equivalents of carboxylate anion per ml of solvent.
9. The composition of claim 8, wherein said acid is a mineral acid, more specifically an acid selected from the group consisting of HCl, HBr, HF, HI,  $H_2SO_4$  and  $HNO_3$  or acetic acid.
10. Method for preparing the stabilized composition according to one or more of claims 1 to 7, comprising the steps of:  
preparing the solvent and treating said solvent to reduce the carboxylate content, and  
dispersing the pharmaceutical compound in said solvent.
11. Method of claim 10, wherein said treating step comprises contacting said solvent with aluminum oxide to reduce the carboxylate anion content of the solvent.
12. Method of claim 10, wherein said treating step comprises adding an acid in an amount to provide at least  $5.6 \times 10^{-6}$ , more specifically about  $5.6 \times 10^{-6}$  to  $8.4 \times 10^{-6}$  grams of  $H^+$  per ml of said solvent.
13. Method of claim 12, wherein said acid is a mineral acid, more specifically an acid selected from the group consisting of HCl, HBr, HF, HI,  $H_2SO_4$  and  $HNO_3$  or acetic acid.

## Patentansprüche

1. Stabilisierte Zusammensetzung umfassend mindestens eine pharmazeutische Verbindung ausgewählt aus der Gruppe bestehend aus Paclitaxel, Camptothecin und Derivaten davon; und  
 5 ein Lösemittel, das fähig ist, die pharmazeutische Verbindung zu dispergieren oder zu solubilisieren, wobei das Lösemittel eine wirksame Menge eines nichtionischen Solubilisierungsmittels enthält mit einem Gehalt an Carboxylatanion, der ausreichend niedrig ist, daß ein katalysierter Abbau der pharmazeutischen Verbindung verhindert ist,  
 10 vorausgesetzt, daß Verbindungen, die Paclitaxel, polyethoxyliertes Castoröl und Ethanol enthalten, eingestellt auf einen pH von weniger als 8,1 durch Zugabe einer anderen Säure als HBr, HF und HI ausgeschlossen sind.
2. Stabilisierte Zusammensetzung umfassend mindestens eine pharmazeutische Verbindung ausgewählt aus der Gruppe bestehend aus Paclitaxel, Camptothecin und Derivaten davon; und  
 15 ein Lösemittel, das fähig ist, die pharmazeutische Verbindung zu dispergieren oder zu solubilisieren, wobei das Lösemittel eine wirksame Menge eines nichtionischen Solubilisierungsmittels enthält mit einem Gehalt an Carboxylatanion, der ausreichend niedrig ist, daß ein katalysierter Abbau der pharmazeutischen Verbindung verhindert ist,  
 20 worin vor dem Dispergieren der pharmazeutischen Verbindung in dem Lösemittel, das Lösemittel behandelt wurde, um den Gehalt an Carboxylatanion zu reduzieren, wobei die Behandlung in Kontakt bringen des Lösemittels mit Aluminiumoxid umfaßt.
3. Zusammensetzung nach Anspruch 1 oder 2, worin das Lösemittel einen Gehalt an Carboxylatanion aufweist von ungefähr  $0,6 \times 10^{-6}$  Grammäquivalenten Carboxylatanion pro ml Lösemittel.
- 25 4. Zusammensetzung nach Anspruch 1, 2 oder 3, worin das Lösemittel ein Kondensationsprodukt von Alkylenoxid und einem Lipid umfaßt, insbesondere polyoxyethyliertes Castoröl oder polyoxyethylierte 12-Hydroxystearinsäure.
5. Zusammensetzung nach Anspruch 4, worin das polyoxyethylierte Lipidkondensationsprodukt hergestellt ist aus Castoröl und 20 bis 40 Mol Ethylenoxid pro Mol Castoröl.
- 30 6. Zusammensetzung nach einem der Ansprüche 1 bis 5, worin das Lösemittel ferner einen Alkohol umfaßt, insbesondere einen Alkohol ausgewählt aus der Gruppe bestehend aus Ethanol und Polyethylenglykol.
7. Zusammensetzung nach Anspruch 6, worin das Lösemittel eine Mischung aus Ethylalkohol und polyoxyethyliertem Castoröl in einem Volumenverhältnis von ungefähr 50 : 50 ist.
- 35 8. Zusammensetzung nach einem der Ansprüche 1 bis 7, worin das Lösemittel eine wirksame Menge einer Säure enthält, so daß ein Gehalt an Carboxylatanion von weniger als oder gleich ungefähr  $0,6 \times 10^{-6}$  Grammäquivalenten Carboxylatanion pro ml Lösemittel eingehalten ist.
- 40 9. Zusammensetzung nach Anspruch 8, worin die Säure eine Mineralsäure ist, insbesondere eine Säure ausgewählt aus der Gruppe bestehend aus HCl, HBr, HF, HI,  $H_2SO_4$  und  $HNO_3$  oder Essigsäure.
- 45 10. Verfahren zur Herstellung der stabilisierten Zusammensetzung nach einem der Ansprüche 1 bis 7, umfassend die Schritte:  
 Herstellen des Lösemittels und Behandeln des Lösemittels, um den Carboxylatgehalt zu reduzieren, und Dispergieren der pharmazeutischen Verbindung in dem Lösemittel.
- 50 11. Verfahren nach Anspruch 10, worin der Behandlungsschritt in Kontakt bringen des Lösemittels mit Aluminiumoxid umfaßt, so daß der Gehalt des Lösemittels an Carboxylatanion reduziert wird.
12. Verfahren nach Anspruch 10, worin der Behandlungsschritt Zugabe einer Säure in einer Menge umfaßt, so daß mindestens  $5,6 \times 10^{-6}$ , insbesondere ungefähr  $5,6 \times 10^{-6}$  bis  $8,4 \times 10^{-6}$  Gramm  $H^+$  pro ml des Lösemittel erreicht werden.
- 55 13. Verfahren nach Anspruch 12, worin die Säure eine Mineralsäure ist, insbesondere eine Säure ausgewählt aus der Gruppe bestehend aus HCl, HBr, HF, HI,  $H_2SO_4$  und  $HNO_3$  oder Essigsäure.

## Revendications

1. Composition stabilisée comprenant au moins un composé pharmaceutique sélectionné dans le groupe consistant en paclitaxel, camptothécine et leurs dérivés; et  
 un solvant capable de disperser ou de solubiliser ledit composé pharmaceutique, ledit solvant comprenant une quantité efficace d'un agent solubilisant non ionique ayant une teneur en anion carboxylate suffisamment faible pour empêcher la dégradation catalysée dudit composé pharmaceutique,  
 à condition que les compositions comprenant le paclitaxel, de l'huile de ricin polyéthoxylée et de l'éthanol ajusté à un pH inférieur à 8,1 par addition d'un acide autre que HBr, HF et HI, soient exclues.
2. Composition stabilisée comprenant au moins un composé pharmaceutique sélectionné dans le groupe consistant en paclitaxel, camptothécine et leurs dérivés; et  
 un solvant capable de disperser ou de solubiliser ledit composé pharmaceutique, ledit solvant comprenant une quantité efficace d'un agent solubilisant non ionique ayant une teneur en anion carboxylate suffisamment faible pour empêcher la dégradation catalysée dudit composé pharmaceutique où, avant dispersion dudit composé pharmaceutique dans ledit solvant, ledit solvant a été traité pour réduire la teneur en anion carboxylate, ledit traitement comprenant la mise en contact dudit solvant avec de l'oxyde d'aluminium.
3. Composition de la revendication 1 ou 2, où ledit solvant a une teneur en anion carboxylate d'environ  $0,6 \times 10^{-6}$  équivalent gramme d'anion carboxylate par ml du solvant.
4. Composition de la revendication 1, 2 ou 3, où ledit solvant comprend un produit de condensation d'oxyde d'alkylène et d'un lipide, plus particulièrement d'huile de ricin polyéthoxylée ou d'acide 12-hydroxy stéarique polyéthylé.
5. Composition de la revendication 4, où ledit produit de condensation lipidique polyoxyéthylé est fait d'huile de ricin et de 20 à 40 moles d'oxyde d'éthylène par mole d'huile de ricin.
6. Composition selon l'une des revendications 1 à 5, où ledit solvant contient de plus un alcool, plus particulièrement un alcool sélectionné dans le groupe consistant en éthanol et polyéthylène glycol.
7. Composition de la revendication 6, où ledit solvant est un mélange d'alcool éthylique et de ladite huile de ricin polyoxyéthylée à un rapport d'environ 50:50 en volume.
8. Composition selon l'une des revendications 1 à 7, où ledit solvant contient une quantité efficace d'un acide pour maintenir la teneur en anion carboxylate à moins de ou égale à environ  $0,6 \times 10^{-6}$  équivalent gramme d'anion carboxylate par ml du solvant.
9. Composition de la revendication 8, où ledit acide est un acide minéral, plus particulièrement un acide sélectionné dans le groupe consistant en HCl, HBr, HF, HI,  $H_2SO_4$  et  $HNO_3$  ou l'acide acétique.
10. Méthode de préparation de la composition stabilisée selon une ou plusieurs des revendications 1 à 7, comprenant les étapes de :  
 préparer le solvant et traiter ledit solvant pour réduire la teneur en carboxylate, et  
 disperser le composé pharmaceutique dans ledit solvant.
11. Méthode de la revendication 10, où ladite étape de traitement comprend la mise en contact dudit solvant avec de l'oxyde d'aluminium pour réduire la teneur en anion carboxylate du solvant.
12. Méthode de la revendication 10, où ladite étape de traitement comprend l'addition d'un acide en une quantité pour qu'il y ait au moins  $5,6 \times 10^{-6}$ , plus particulièrement environ  $5,6 \times 10^{-6}$  à  $8,4 \times 10^{-6}$  gramme de  $H^+$  par ml dudit solvant.
13. Méthode de la revendication 12, où ledit acide est un acide minéral, plus particulièrement un acide sélectionné dans le groupe consistant en HCl, HBr, HF, HI,  $H_2SO_4$  et  $HNO_3$  ou l'acide acétique.